

Development and Validation of the Japanese version of the Person-Centered Primary Care Measure (Research in progress)

Makoto Kaneko^{a,b}, Tadao Okada^c, Takuya Aoki^d, Machiko Inoue^b, Takamasa Watanabe^e, Masato Matsushima^d

a. Primary Care Research Unit, Graduate School of Health Data Science, Yokohama City University b. Department of Family and Community Medicine, Hamamatsu University School of Medicine c. Tessayukai Kameda Family Clinic Tateyama d. Division of Clinical Epidemiology, Jikei University School of Medicine e. Centre for Family Medicine Development, Japanese Health and Welfare Co-operative Federation

Background		Objectives		Criterion validity	Using JPCAT	
<ul style="list-style-type: none"> • Patient Experience is an important quality indicator. • In Japan, the Japanese version of Primary Care Assessment Tool (JPCAT) has been widely used since 2016.¹ • Person-Centered Primary Care Measure (PCPCM) by Etz et al. 2019, can assess a wider domain of primary care and compare the results with other countries directly.² 		This study aims to develop and validate the Japanese version of the PCPCM.		Convergent and Discriminant validity	Examine an association between PCPCM and vaccine uptake	
		Methods		<u>Study Design</u> A cross-sectional survey <u>Population studied</u> 1,000 participants selected by population-weighted random sampling from the Basic Resident Registration System in Japan <u>Outcome Measures</u> Reliability Internal consistency • Cronbach's α • Item-total correlations Validity Structural validity Exploratory factor analysis: eigenvalue, scree plot		Current progress <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; background-color: #76b82a; padding: 5px;">Translation - back translation</div> <div style="border: 1px dashed black; padding: 5px;">Cognitive debriefing</div> <div style="border: 1px dashed black; padding: 5px;">Validity Reliability</div> </div> <div style="border: 1px solid black; background-color: #76b82a; width: 100px; height: 20px; margin: 10px auto;"></div> <div style="border: 1px dashed black; width: 100%; height: 20px; margin: 10px auto;"></div>
PCPCM 11 domain (*JPCATdomain)				Your suggestion/advice wanted ! <ul style="list-style-type: none"> • Is sampling strategy appropriate? General population? Clinical sampling? Mixed? • Can we evaluate reliability and validity in this way appropriately? 		
Advocacy	Accessibility*					
Comprehensiveness*	Community context*					
Coordination*	Continuity*					
Family context	Integration					
Goal-oriented care	Health promotion					
	Relationship					

