

References

- Adekanattu, P., Sholle, E. T., DeFerio, J., Pathak, J., Johnson, S. B., & Campion, J., Thomas R. (2018). Ascertaining depression severity by extracting patient health questionnaire 9 (PHQ-9) scores from clinical notes. *AMIA . Annual Symposium Proceedings. AMIA Symposium, 2018*, 147-156.
- Bhattacharjee, S., Goldstone, L., Vadiiei, N., Lee, J. K., & Burke, W. J. (2018). Depression screening patterns, predictors, and trends among adults without a depression diagnosis in ambulatory settings in the United States. *Psychiatric Services, 69*(10), 1098-1100.
doi: 10.1176/appi.ps.201700439
- Thompson, H., Faig, W., Gupta, N., Lahey, R., Golden, R., Pollack, M., & Karnik, N. (2019). Collaborative care for depression of adults and adolescents: Measuring the effectiveness of screening and treatment uptake. *Psychiatric Services, 70*(7), 604-607.
doi:10.1176/appi.ps 201800257
- United States Preventive Services Task Force. (2016). Recommendations for primary care practice. <https://www.uspreventiveservicestaskforce.org/>
Link: <https://uspreventiveservicestaskforce.org/uspstf/recommendation/depression-in-adults-screening>

